

Verified Estimation of Taylor Coefficients and Taylor Remainder Series of Analytic Functions

Markus Neher

Institut für Angewandte Mathematik, Universität Karlsruhe
D-76128 Karlsruhe, Germany
markus.neher@math.uni-karlsruhe.de

1 Introduction

Let $y = \sum_{j=0}^{\infty} b_j z^j$ be the analytic solution of a problem $F(y) = 0$ that depends on analytic functions f_1, \dots, f_n and that can be solved by recurrent computation of the Taylor coefficients b_j of y . Only a finite number of the b_j can be calculated in practical computations. Then a finite sum $\sum_{j=0}^p b_j z^j$ only yields an approximation to $y(z)$.

In some cases, however, it is possible to determine a geometric series (or some derivative of a geometric series) that serves as a bound for the remainder series of y , provided that the remainder series of the functions f_1, \dots, f_n can also be estimated by geometric series (or derivatives of geometric series). In [4], such an error analysis was used for the validated solution of linear ODEs.

A prerequisite of this general method is the computation of bounds for the Taylor coefficients of arbitrary order of a given analytic function f . The subject of this talk is the validated solution of the latter problem.

2 Estimates for Taylor Coefficients

In the following, let $f(z) = \sum_{j=0}^{\infty} a_j z^j$ be analytic in B and bounded on C , where B is the complex disc $\{z : |z| < r\}$ and C the circle $\{z : |z| = r\}$, for some $r > 0$. A well known bound for the Taylor coefficients of f is Cauchy's estimate $M(r)$:

$$|a_j| \leq \frac{M(r)}{r^j}, \quad M(r) := \max_{|z|=r} |f(z)|, \quad j \in \mathbb{N}_0.$$

Unfortunately, Cauchy's estimate is sometimes very pessimistic. To obtain better bounds, two modifications of Cauchy's estimate were proposed in [5]. The first uses a Taylor polynomial to approximate f :

Theorem 1 Let f be analytic in B and bounded on C . Furthermore, let $T_l(z)$ denote the Taylor polynomial of order l to f . Then

$$|a_j| \leq \frac{N(r, l)}{r^j} \quad \text{for } j > l, \quad \text{where } N(r, l) := \max_{|z|=r} |f(z) - T_l(z)|.$$

Cauchy's estimate can also be improved using the derivatives of f :

Theorem 2 Let f be analytic in B and let $f^{(m)}$ (the m -th derivative of f) be bounded on C . Furthermore, let $P(j, m) := (j+1) \cdots (j+m)$, $P(j, 0) := 1$ for $m \in \mathbb{N}$, $j \in \mathbb{N}_0$. Then

$$|a_j| \leq \frac{U(r, m)r^m}{P(j-m, m)r^j} \quad \text{for } j \geq m, \quad \text{where } U(r, m) := \max_{|z|=r} |f^{(m)}(z)|.$$

3 Estimates for Taylor Remainder Series

The estimation of the remainder series $R_p := \sum_{j=p+1}^{\infty} a_j z^j$ is obtained by addition of the above estimates of the Taylor coefficients. Using Theorem 1, at some point z with $|z| = \omega r$, $\omega \in (0, 1)$, for arbitrary $p \geq l$ we have

$$|R_p(z)| \leq \sum_{j=p+1}^{\infty} N(r, l) \omega^j = N(r, l) \frac{\omega^{p+1}}{1-\omega},$$

whereas for $p \geq m-1$ we have

$$|R_p(z)| \leq \sum_{j=p+1}^{\infty} \frac{U(r, m)r^m}{P(j-m, m)} \omega^j$$

by Theorem 2. For $p = m-1$ the latter sum is derived by repeated integration of $\frac{1}{1-\omega}$. For $p \gg m$ the resulting formula suffers from severe cancellation. In this case, the estimate

$$R_p \leq \frac{U(r, m)r^m}{P(p+1-m, m)} \frac{\omega^{p+1}}{1-\omega}$$

is better suited for practical calculations.

In the situation that was mentioned in the introduction, it is usually not known in advance which order p of the Taylor polynomial T_p is required for sufficient accuracy of the approximation of the unknown solution y . Because it is very expensive to recalculate the bounds N or U for different values of l or m , respectively, it is better to guess sufficiently large values for l or m a priori and to calculate N or U only once. Only if $p = l$ or $p = m$ are not sufficient for a good approximation of y , then p is augmented iteratively, but R_p is still calculated for the same values of l or m (note that $R_p \rightarrow 0$ for $p \rightarrow \infty$ independently of l or m).

4 Implementation

Using interval arithmetic [1] on the computer, the validated computation of the above estimates is possible for analytic compositions of rational functions and of those complex standard functions that are available on the computer (like e^z , $\sin z$, $\text{Log } z$, \dots). The real and imaginary parts of many standard functions can be expressed as compositions of real standard functions. Such compositions have been utilized in [2] for the construction of complex interval standard functions that enclose the respective ranges over complex intervals. These inclusion functions and well known methods for rigorous global optimization [3] are used in the practical calculation of the estimates $M(r)$, $N(r, l)$, or $U(r, m)$.

5 ACETAF Software

The above algorithms for the computation of guaranteed upper bounds for the Taylor coefficients and remainder series of analytic functions have been implemented in a C-XSC program called ACETAF. The program also contains routines for the validated automatic computation of derivatives of complex analytic functions and for the check of analyticity of user-defined functions in circles in the complex plane.

ACETAF is distributed under the terms of the GNU General Public License and is available at the following site:

<http://www.uni-karlsruhe.de/~Markus.Neher/acetaf.html>

References

- [1] G. Alefeld and J. Herzberger, *Introduction to interval computations*, Academic Press, New York, 1983.
- [2] K. Braune and W. Krämer, “High-accuracy standard functions for real and complex intervals”, In: E. Kaucher, U. Kulisch, and Ch. Ullrich (eds), *Computerarithmetic: Scientific computation and programming languages*, Teubner, Stuttgart, 1987, pp. 81–114.
- [3] R. B. Kearfott, *Rigorous global search: Continuous problems*, Kluwer, Dordrecht, 1996.
- [4] M. Neher, “Geometric series bounds for the local errors of Taylor methods for linear n -th order ODEs”, In: G. Alefeld, J. Rohn, S. Rump, and T. Yamamoto (eds), *Symbolic Algebraic Methods and Verification Methods*, Springer, Wien, 2001, pp. 183–193.
- [5] M. Neher, “Validated bounds for Taylor coefficients of analytic functions”, *Reliable Computing*, 2001, Vol. 7, pp. 307–319.