A New Differential Formalism for Interval-Valued Functions and Its Potential Use in Detecting 1-D Landscape Features

Vladik Kreinovich[1], Hung T. Nguyen[1],
Graçaliz Pereira Dimuro[3],
Antônio Carlos da Rocha Costa[3],
and Benjamin Rene Callejas Bedregal[4]

[1]Department of Computer Science
University of Texas at El Paso
El Paso, TX 79968, USA
vladik@cs.utep.edu

[2]Department of Mathematical Sciences
New Mexico State University
Las Cruces, NM 88003, USA
hunguyen@nmsu.edu

[3]Escola de Informática, Universidade Católica de Pelotas
Rua Felix da Cunha 412, 96010-000, Pelotas, Brazil
liz@atlas.ucpel.tche.br, rocha@atlas.ucpel.tche.br

[4]Laboratorio de Logica e Inteligencia Computacional–LabLIC
Departamento de Informática e Matemática Aplicada–DIMAp
Universidade Federal do Rio Grande do Norte–UFRN
Campus Universitario, Lagoa Nova, Natal-RN-Brasil
bedregal@dimap.ufrn.br

Abstract

In many practical problems, it is important to know the slope (derivative) dy/dx of one quantity y with respect to some other quantity x. For example, different 1-D landscape features can be characterized by differ-

ent values of the derivative dy/dx, where y is an altitude, and x is a horizontal coordinate. In practice, we often know the values of $y(x)$ for different x with interval uncertainty. How can we then find the set of possible values of the slope? In this paper, we formulate this problem of differentiating interval-values functions in precise terms, and we describe an (asymptotically) optimal algorithm for computing the corresponding derivative.

1 Introduction

In many real-life problems, we want to know the values of the derivatives. In many areas of science and engineering, we are interested in slopes. For example, a 1-D landscape is described as a dependence of the altitude y on the coordinate x; different 1-D landscape features are defined by different values of the slope dy/dx of this dependence: low values of this slope correspond to a plain, high values to steep mountains, and medium values to a hilly terrain.

Interval uncertainty. In the ideal situation, when we know the exact values of $y(x)$ for every x, we can simply differentiate the corresponding dependence. In practice, however, the information on y comes from measurements, and measurements are never exact. E.g., in the landscape example, we measure the altitudes $y_1, \ldots, y_n$ at different points $x_1 < \ldots < x_n$. Since the measurements are not exact, the measured values $\widetilde{y}_i$ are, in general, slightly different from the the (unknown) actual altitudes y_i.

For measuring instruments, we usually have an upper bound Δ_i on the measurement error $\Delta y_i \stackrel{\text{def}}{=} \widetilde{y}_i - y_i$ provided by the manufacturer of this instrument: $|\Delta y_i| \le \Delta_i$. Thus, after the measurement, the only information that we have about the actual (unknown) value y_i is that this value belongs to the interval $\mathbf{y}_i = [\underline{y}_i, \overline{y}_i]$, where $\underline{y}_i \stackrel{\text{def}}{=} \widetilde{y}_i - \Delta_i$ and $\overline{y}_i \stackrel{\text{def}}{=} \widetilde{y}_i + \Delta_i$ (for a more detailed description of interval uncertainty, see, e.g., [6, 7, 8, 11]).

Thus, the only information that we have about the actual dependence $y = f(x)$ of y on x is that the (unknown) function $f(x)$ belongs to the class

$$F \stackrel{\text{def}}{=} \{f(x) \mid f(x_i) \in \mathbf{y}_i \text{ for all } i = 1, \ldots, n\}. \tag{1}$$

We also know that the (unknown) function $f(x)$ is smooth (differentiable) – because otherwise, the notion of a slope does not make sense.

In many practical applications, the derivative has a physical meaning, and this meaning implies that it is itself a continuous (or even differentiable) function. For example, when we monitor the locations y_i of a particle at different moments of time x_i, then the derivative dy/dx is a velocity; when we monitor the values y_i of the velocity, then the derivative dy/dx is the acceleration, etc. Thus, we can assume that the function f is continuously differentiable.

How can we determine the slopes under such interval uncertainty?

Toward a formal definition. Let us assume that we look for areas where the slope takes a given value s. In a simplified example, we monitor the location y_i of a car on a highway at different moments of time, and we want to find out where the car was driving at the maximal allowed speed s (or, alternatively, where it was driving at an excessive speed s).

Since we only know the values of the unknown function $f(x)$ at finitely many points $x_1 < \ldots < x_n$, it is always possible that the derivative of the (unknown) function $f(x)$ attains the desired value s at some point between x_i and x_{i+1}. For example, if we are checking for the areas where the car was overspeeding, it is always possible that the car was going very fast when no one was looking (i.e., in between x_i and x_{i+1}), for a short period of time, just for fun, so that the overall traveled distance was not affected.

In other words, for every interval $[a, b]$ $(a < b)$, it is always possible to have a function f within the class F (defined by the formula (1)) for which $f'(x) = s$ for some $s \in [a, b]$.

What we are really interested in is not whether it is *possible* that somewhere, the slope is equal to s (it is always possible), but whether the data *imply* that somewhere, the slope was indeed equal to x. This "implies" means that whatever function $f =\in F$ we take, there always is a point $x \in [a, b]$ for which $f'(x) = s$ (this point may be different for different functions $f \in F$).

In other words, we say that the slope is guaranteed to attain a given value s somewhere on a given interval $[a, b]$ if for every function $f \in F$, the range $f'([a, b])$ of its derivative $f'(x)$ contains the value s. In mathematical terms, this means that the value s belongs to the *intersection* of the ranges $f'([a, b])$ corresponding to all $f \in F$.

This intersection thus describes the "range of the derivative" of the interval-valued function F on the given are $[a, b]$. In other words, we arrive at the following definitions.

2 Precise Formulation of the Problem

Definition 1. *By an* interval-valued function F*, we mean a finite sequence of pairs* $\langle x_i, \mathbf{y}_i \rangle$ *(*$i = 1, 2, \ldots, n$*), where for each* i*,* x_i *is a real number,* $\mathbf{y}_i$ *is a non-degenerate interval, and* $x_1 < x_2 < \ldots < x_n$*.*

Definition 2. *We say that a function* $f : R \to R$ *from reals to reals* belongs *to an interval-valued function* $F = \{\langle x_1, \mathbf{y}_1 \rangle, \ldots, \langle x_n, \mathbf{y}_n \rangle\}$ *if* $f(x)$ *is continuously differentiable and for every* i *from 1 to* n*, we have* $f(x_i) \in \mathbf{y}_i$*.*

Definition 3. *Let F be an interval-valued function, and let $[a,b]$ be an interval. By a* derivative $F'([a,b])$*, we mean the intersection*

$$F'([a,b]) \stackrel{\text{def}}{=} \bigcap_{f \in F} f'([a,b]),$$

where $f'(x)$ denotes the derivative of a differentiable function $f(x)$, and $f'([a,b]) \stackrel{\text{def}}{=} \{f'(x) \mid x \in [a,b]\}$ is the range of the derivative $f'(x)$ over the interval $[a,b]$.

Comment. The notation $F'([a,b])$ looks like the notation of a range for a real-valued function, but it is not a range: in contrast to range, if an interval is narrow enough, we can have $F'([a,b]) = \emptyset$ (see examples below).

This newly defined derivative does share some properties of the range: e.g., since the range is *inclusion-monotonic* – in the sense that $[a,b] \subseteq [c,d]$ implies $f'([a,b]) \subseteq f'([c,d])$ – we conclude that $[a,b] \subseteq [c,d]$ implies $F'([a,b]) \subseteq F'([c,d])$. Thus, if the union $A \cup B$ of two intervals is also an interval, we have $F'(A \cup B) \supseteq F'(A) \cup F'(B)$.

Formulation of the problem. How can we compute the derivative of an interval-valued function? The above definition, if taken literally, requires that we consider all (infinitely many) functions $f \in F$ – which is computationally excessive. Thus, we must find an efficient algorithm for computing this derivative. This is what we will do in this paper.

We will try our best to make sure that these algorithms are not simply tricks, that the ideas behind these algorithms are clear and understandable. Therefore, instead of simply presenting the final algorithm, we will, instead, present our reasoning in a series of auxiliary results that eventually leads to the asymptotically optimal algorithms for computing the desired derivative $F'([a,b]$.

Previous work. In our research, we were guided by results from two related research directions:

- first, by different definitions of differentiation of an interval function that have been proposed by interval computations community [2, 7, 10, 12, 13, 15, 16, 17, 18, 19]; the main difference from our problem is that most of these papers assume that we have intervals $\mathbf{y}$ for all x, while we consider a more realistic situation when the interval bounds on $f(x)$ are only known for finitely many values $x_1, \ldots, x_n$;
- second, by a paper [20] in which an algorithm was developed to check for local maxima and minima of an interval-valued function f; this result has been applied to detecting geological areas [1, 3, 4] and to financial analysis [5]; this result can be viewed as detecting the areas where the derivative is equal to 0 – and, in this sense, as a particular case of our current problem.

3 First Auxiliary Result: Checking Monotonicity

Definition 4. *We say that a function $f(x)$ is* strongly increasing *if $f'(x) > 0$ for all x.*

Comment. Every strongly increasing function is strictly increasing, but the inverse is not necessarily true: the function $f(x) = x^3$ is strictly increasing but not strongly increasing.

Proposition 1. *For every interval function F, the existence of a strongly increasing function $f \in F$ with $f'(x) > 0$ is equivalent to*

$$\underline{y}_i < \overline{y}_j \text{ for all } i < j. \tag{2}$$

Proof. If $f \in F$ and $f(x)$ is strongly increasing, then it is also strictly increasing hence for every $i < j$, the inequality $x_i < x_j$ implies that $f(x_i) < f(x_j)$. Since $f \in F$, we have $f(x_i) \in \mathbf{y}_i = [\underline{y}_i, \overline{y}_i]$ and $f(x_j) \in \mathbf{y}_i = [\underline{y}_j, \overline{y}_j]$. Thus, from $\underline{y}_i \le f(x_i) < f(x_j) \le \overline{y}_j$, we conclude that $\underline{y}_i < \overline{y}_j$, which is exactly the inequality (2).

Vice versa, let us assume that the inequalities (2) are satisfied, and let us design the corresponding strictly increasing function $f \in F$. We will first build a piece-wise linear strictly increasing function $f_0(x)$ for which $f_0(x_i) \in \mathbf{y}_i$, and then we will show how to modify $f_0(x)$ into a continuously differentiable strongly increasing function $f \in F$.

According to the inequalities (2), all the differences $\overline{y}_j - \underline{y}_i$ $(i < j)$ are positive. Since all intervals are non-degenerate, the differences $\overline{y}_i - \underline{y}_i$ are also positive. Let us denote the smallest of these positive numbers by Δ. For every i, let us denote

$$y_i \stackrel{\text{def}}{=} \max(\underline{y}_1, \ldots, \underline{y}_i) + \frac{i}{2n} \cdot \Delta. \tag{3}$$

We will then design $f_0(x)$ as a piece-wise linear function for which $f_0(x_i) = y_i$. To show that $f_0(x)$ is the desired piece-wise linear function, we must show that for every i, $y_i \in \mathbf{y}_i$, and that this function is strictly increasing, i.e., that $i < j$ implies $y_i < y_j$.

That $i < j$ implies $y_i < y_j$ is clear: the first (maximum) term in the formula (3) can only increase (or stay the same) when we replace i by j, and the second term increases. Thus, it is sufficient to prove that $y_i \in \mathbf{y}_i = [\underline{y}_i, \overline{y}_i]$, i.e., that $\underline{y}_i \le y_i$ and $y_i \le \overline{y}_i$. We will actually prove a stronger statement: that $\underline{y}_i < y_i$ and $y_i < \overline{y}_i$.

The first inequality $\underline{y}_i < y_i$ follows directly from the formula (3): by definition of a maximum, $\max(\underline{y}_1, \ldots, \underline{y}_i) \geq \underline{y}_i$, and when we add a positive number to this maximum, the result only increases. So, y_i is actually larger than $\underline{y}_i$.

Let us now prove that $y_i < \overline{y}_i$. Indeed, by definition of Δ, for all $k \leq i$, we have $\underline{y}_k + \Delta \leq \overline{y}_i$, hence (since $(i/2n) \cdot \Delta < \Delta$) $\underline{y}_k + (i/2n)\Delta < \overline{y}_i$. Thus, y_i – which is the largest of the values $\underline{y}_k + (i/2n)\Delta$ – is also smaller than $\overline{y}_i$. So, the desired $f_0(x)$ is designed.

Let us now show how to build the corresponding continuously differentiable function $f(x)$. For the piece-wise linear function $f_0(x)$, the first derivative $f_0'(x)$ is piece-wise constant; since the function $f_0(x)$ is strictly increasing, the values $f_0'(x)$ are all positive. Around each discontinuity point x_i, replace the abrupt transition with a linear one; as we integrate the resulting function, we get a new function $f(x)$ that is continuously differentiable and – since the new values of the derivative are still everywhere positive – strongly increasing. When the replacement is fast enough, the change in the value $f(x_i)$ is so small that $f(x_i)$ is still inside the desired interval $\mathbf{y}_i$. The proposition is proven.

Similarly, we can prove the following results:

Definition 5. *We say that a function $f(x)$ is* strongly decreasing *if $f'(x) < 0$ for all x.*

Proposition 2. *For every interval function F, the existence of a strongly decreasing function $f \in F$ is equivalent to*

$$\overline{y}_i > \underline{y}_j \text{ for all } i < j. \tag{4}$$

Proposition 3. *For every interval function F and for every interval $[a, b]$, the existence of a function $f \in F$ that is strongly increasing on the interval $[a, b]$ is equivalent to*

$$\underline{y}_i < \overline{y}_j \text{ for all } i < j \text{ for which } x_i, x_j \in [a, b]. \tag{5}$$

Proposition 4. *For every interval function F and for every interval $[a, b]$, the existence of a function $f \in F$ that is strongly decreasing on the interval $[a, b]$ is equivalent to*

$$\overline{y}_i > \underline{y}_j \text{ for all } i < j \text{ for which } x_i, x_j \in [a, b]. \tag{6}$$

4 Second Auxiliary Result: When $0 \in F'([a,b])$

Proposition 5. *For every interval function F and for every interval $[a,b]$, $0 \in F'([a,b])$ if and only if neither conditions (5) not conditions (6) are satisfied.*

Proof. Let us first show that if either the conditions (5) or the conditions (6) are satisfied, then $0 \notin F'([a,b])$.

Indeed, according to Proposition 3, if the conditions (5) are satisfied, then there exists a function $f \in F$ that is strongly increasing on $[a,b]$. For this function, $f'(x) > 0$ for all $x \in [a,b]$; therefore, $f'([a,b]) \subseteq (0,\infty)$. Since $F'([a,b])$ is defined as the intersection of such range sets, we have $F'([a,b]) \subseteq f'([a,b]) \subseteq (0,\infty)$ hence $0 \notin F'([a,b])$.

Similarly, if the conditions (6) are not satisfied, then $0 \notin F'([a,b])$.

Vice versa, let us assume that neither the conditions (5) nor the conditions (6) are satisfied, and let us show that then $0 \in F'([a,b])$. Indeed, let $f \in F$ be an arbitrary function from the class F. Since the conditions (5) are not satisfied, the function $f(x)$ cannot be strongly increasing; therefore, there must be a point $x_1 \in [a,b]$ for which $f'(x_1) \le 0$. Similarly, since the conditions (6) are not satisfied, the function $f(x)$ cannot be strongly decreasing; therefore, there must be a point $x_2 \in [a,b]$ for which $f'(x_2) \ge 0$.

Since the function $f(x)$ is continuously differentiable, the continuous derivative $f'(x)$ must attain the 0 value somewhere on the interval $[x_1, x_2] \subseteq [a,b]$. In other words, $0 \in f'([a,b])$ for all $f \in F$. Thus, 0 belongs to intersection $F'([a,b])$ of all possible ranges $f'([a,b])$. The proposition is proven.

5 Third Auxiliary Result and Final Description of $F'([a,b])$

Definition 6. *Let*

$$F = \{\langle x_1, \mathbf{y}_1\rangle, \ldots, \langle x_n, \mathbf{y}_n\rangle\}$$

be an interval function, and let v be a real number. Then, we define a new interval function $F - v \cdot x$ as follows:

$$F - v \cdot x = \{\langle x_1, \mathbf{y}_1 - v \cdot x_1\rangle, \ldots, \langle x_n, \mathbf{y}_n - v \cdot x_n\rangle\},$$

where, for an interval $\mathbf{y} = [\underline{y}, \overline{y}]$ and for a real number c, the difference $\mathbf{y} - c$ is defined as $[\underline{y} - c, \overline{y} - c]$.

It is easy to prove the following auxiliary result:

Proposition 6. *For every interval function F and for every interval $[a,b]$, $v \in F'([a,b])$ if and only if $0 \in (F - v \cdot x)'([a,b])$.*

This results leads to the following description of the derivative $F'([a,b])$:

Proposition 7. *For every interval function F and for every interval $[a,b]$, let i_0 and j_0 be the first and the last index of the values x_i inside $[a,b]$. Then $F'([a,b]) = [\underline{F}_{i_0,j_0}, \overline{F}_{i_0 j_0}]$, where*

$$\underline{F}_{i_0 j_0} \stackrel{\text{def}}{=} \min_{i_0 \le i < j \le j_0} \overline{\Delta}_{ij}, \tag{7}$$

$$\overline{F}_{i_0 j_0} \stackrel{\text{def}}{=} \max_{i_0 \le i < j \le j_0} \underline{\Delta}_{ij}, \tag{8}$$

$$\underline{\Delta}_{ij} \stackrel{\text{def}}{=} \frac{\underline{y}_i - \overline{y}_j}{x_j - x_i}, \tag{9}$$

$$\overline{\Delta}_{ij} \stackrel{\text{def}}{=} \frac{\overline{y}_i - \underline{y}_j}{x_j - x_i}, \tag{10}$$

and when $p > q$, the interval $[p,q] = \{x \mid p \le x \,\&\, x \le q\}$ is the empty set.

Comment. The above expression is rather intuitively reasonable because the ratios $\underline{\Delta}_{ij}$ and $\overline{\Delta}_{ij}$ are finite differences – natural estimates for the derivatives.

Proof. The fact that conditions (5) are not satisfied means that there exist value $i_0 \le i < j \le j_0$ for which $\underline{y}_i \ge \overline{y}_j$. The fact that the conditions (6) are not satisfied means that there exist values $i_0 \le i' < j' \le j_0$ for which $\overline{y}_{i'} \le \underline{y}_{j'}$.

Similarly, the fact that the conditions (5) and (6) are not satisfied for the interval function $F - v \cdot x$ mean that

$$\exists i, j\, (i_0 \le i < j \le j_0 \,\&\, \underline{y}_i - v \cdot x_i \ge \overline{y}_j - v \cdot x_j) \tag{11}$$

and

$$\exists i', j'\, (i_0 \le i' < j' \le j_0 \,\&\, \overline{y}_{i'} - v \cdot x_{i'} \le \underline{y}_{j'} - v \cdot x_{j'}). \tag{12}$$

The inequality $\underline{y}_i - v \cdot x_i \ge \overline{y}_j - v \cdot x_j$ can be described in the equivalent form $v \cdot (x_j - x_i) \ge \overline{y}_j - \underline{y}_i$, i.e., since $x_i < x_j$, in the form $v \ge \overline{\Delta}_{ij}$. Thus, the existence of i and j as expressed by the formula (11) can be described as the existence of i and j for which v is larger than the corresponding value $\overline{\Delta}_{ij}$, i.e., as

$$v \ge \min_{i_0 \le i < j \le j_0} \overline{\Delta}_{ij}.$$

Similarly, the condition (12) is equivalent to

$$v \le \max_{i_0 \le i < j \le j_0} \underline{\Delta}_{ij}.$$

The proposition is proven.

6 Towards a Faster Algorithm

Proposition 7 provides an explicit formula for computing $F'([a,b])$ for each interval $[a,b]$. For each $[a,b]$, we need to compute $O(n^2)$ values of $\underline{\Delta}_{ij}$ and $\overline{\Delta}_{ij}$.

In problem like locating landscape features, we are not so much interested in knowing whether a given type of landscape exists in a given zone, but rather in locating all types of landscape. In other words, we would like to be able to find the values $F'([a,b])$ for all possible intervals $[a,b]$. According to Proposition 7, it is sufficient to find all the values $F'([x_{i_0}, x_{j_0}])$ for all $i_0, j_0 = 1, \ldots, n$ for which $i_0 < j_0$. There are $n \cdot (n+1)/2 = O(n^2)$ such values. If we use the formula from Proposition 7 – that takes $O(n^2)$ computational steps – to compute each of these $O(n^2)$ values, we will need an overall of $O(n^2) \cdot O(n^2) = O(n^4)$ steps.

For large n – e.g., for $n \approx 10^6$ – we need $n^4 \approx 10^{24}$ computational steps; this is too long for even the fastest computers. Let us show that we can compute the interval derivative faster, actually in $O(n^2)$ time. Since we must return $O(n^2 0$ results, we cannot do it in less than $O(n^2)$ computational steps – so this algorithm is (asymptotically) optimal.

Proposition 8. *There exists an algorithm that, given an interval function $F = \{\langle x_1, \mathbf{y}_1\rangle, \ldots, \langle x_n, \mathbf{y}_n\rangle\}$, computes all possible values of the derivative $F'([a,b])$ in $O(n^2)$ computational steps.*

Proof. At first, we compute $O(n^2)$ values $\underline{\Delta}_{ij}$ and $\overline{\Delta}_{ij}$ by using the formulas (9) and (10); this requires $O(n^2)$ steps.

Let us now show how to compute all n^2 values $\overline{F}_{i_0 j_0}$ in $O(n^2)$ steps.

First, for each i, we sequentially compute the "vertical" maxima $\overline{v}_{ij} \stackrel{\text{def}}{=} \max(\underline{\Delta}_{i,i+1}, \ldots, \underline{\Delta}_{ij})$ corresponding to $j = i+1, i+2, \ldots, n$ as follows: $\overline{v}_{i,i+1} = \underline{\Delta}_{i,i+1}$ and $\overline{v}_{ij} = \max(\overline{v}_{i,j-1}, \underline{\Delta}_{ij})$ for $j > i+1$. For each $i = 1, \ldots, n$, to compute all these values, we need $\le n$ computational steps. Thus, to compute all such values $\overline{v}_{ij}$ for all i and j, we need $\le n \cdot n = O(n^2)$ computational steps.

Then, for every j_0, we sequentially compute the values $\overline{F}_{i_0 j_0}$ for $i_0 = j_0 - 1, j_0 - 2, \ldots, 1$ as follows: $\overline{F}_{j_0-1,j_0} = \underline{v}_{j_0-1,j_0}$ and $\overline{F}_{i_0,j_0} = \max(\overline{F}_{i_0+1,j_0}, \overline{v}_{i_0,j_0})$ (it is easy to see that this formula is indeed correct). For each $j_0 = 1, \ldots, n$, to compute all these values, we need $\le n$ computational steps. Thus, to compute all such values $\overline{F}_{ij}$ for all i_0 and j_0, we need $\le n \cdot n = O(n^2)$ computational steps.

Similarly, by using $\overline{\Delta}_{ij}$ instead of $\underline{\Delta}_{ij}$ and min instead of max, we can compute all n^2 values $\underline{F}_{i_0 j_0}$ in $O(n^2)$ steps. The proposition is proven.

7 This Same Differential Formalism Also Serves an Alternative Definition of Zones

In some practical problems, a zone is defined not by an exact value of the derivative v, but an interval $\mathbf{v} = [\underline{v}, \overline{v}]$ of possible values. In this case, it makes sense to say that an interval $[a, b]$ contains a zone if for every function $f \in F$, there is at least one point $x \in [a, b]$ for which $f'(x) \in \mathbf{v}$. In other words, we say that the interval $[a, b]$ contains a zone of a given type if $f'([a, b]) \cap \mathbf{v} \neq \emptyset$ for all functions $f \in F$.

It turns out that the above notion of a derivative can help us detect such zones as well. Namely, the following statement is true:

Proposition 9. *For every interval function F and for every two intervals $[a, b]$ and $\mathbf{v}$, the following properties are equivalent to each other:*

- *for every function $f \in F$, we have $f'([a, b]) \cap \mathbf{v} \neq \emptyset$;*
- *$\underline{F}_{i_0 j_0} \leq \overline{v}$ and $\overline{F}_{i_0 j_0} \geq \underline{v}$.*

Proof. We will prove the equivalence of the two opposite statements:

- there exists a function $f \in F$ for which $f'([a, b]) \cap \mathbf{v} = \emptyset$;
- $\underline{F}_{i_0 j_0} > \overline{v}$ or $\overline{F}_{i_0 j_0} < \underline{v}$.

Indeed, let us assume that there exists a function $f \in F$ for which $f'([a, b]) \cap \mathbf{v} = \emptyset$. Since every function $f \in F$ is continuously differentiable, its derivative $f'(x)$ is a continuous function, hence the range $f'([a, b])$ is an interval. There are two possible situations when this interval range does not intersect with $\mathbf{v}$:

- either all the values from this range are $> \overline{v}$,
- or all the values from this range are $< \underline{v}$.

In the first case, we have $f'(x) > \overline{v}$ for all $x \in [a, b]$. Therefore, for the function $g(x) \stackrel{\text{def}}{=} f(x) - \overline{v} \cdot x$, we get $g'(x) > 0$ for all x, i.e., the function $g(x)$ is strongly increasing. Since $f \in F$, we have $g \in G \stackrel{\text{def}}{=} F - \overline{v} \cdot x$. Due to Proposition 1, the existence of a strongly increasing function $g \in G$ means that $\underline{y}_i - \overline{v} \cdot x_i < \overline{y}_j - \overline{v} \cdot x_j$ for all $i < j$. This inequality, in its turn, means that $\overline{\Delta}_{ij} > \overline{v}$ for all $i < j$. Thus, $\overline{v}$ is smaller than the smallest of the values $\overline{\Delta}_{ij}$, i.e., smaller than $\underline{F}_{i_0 j_0}$.

Similarly, in the second case, we have $f'(x) < \underline{v}$ for all $x \in [a, b]$, hence $\overline{F}_{i_0 j_0} < \underline{v}$.

Vice versa, let $\underline{F}_{i_0 j_0} > \overline{v}$. By definition of $\underline{F}_{i_0 j_0}$ as the minimum, this means that $\overline{\Delta}_{ij} > \overline{v}$ for all i, j for which $i_0 \leq i < j \leq j_0$. Substituting the definition of $\overline{\Delta}_{ij}$, multiplying both sides of the inequality by a positive term $x_j - x_i$ and

moving terms to another side, we conclude that $\underline{y}_i - \overline{v} \cdot x_i < \overline{y}_j - \overline{v} \cdot x_j$ for all $i < j$. This inequality, in its turn, means that for the interval function $G \stackrel{\text{def}}{=} F - \overline{v} \cdot x$, formula (2) holds and thus, due to Proposition 1, there exist a strongly monotonic function $g \in G$ for which $g'(x) > 0$ for all x. Then, for the function $f(x) \stackrel{\text{def}}{=} g(x) + \overline{v} \cdot x$, we have $f \in F$ and $f'(x) = g'(x) + \overline{v} > \overline{v}$ for all x – hence, $f'([a, b]) \cap \mathbf{v} = \emptyset$.

Similarly, if $\overline{F}_{i_0 j_0} < \underline{v}$, there exists a function $f \in F$ for which $f'([a, b]) \cap \mathbf{v} = \emptyset$. The proposition is proven.

8 Open Problems

What if we take into consideration uncertainty in measuring x? In the above text, we took into consideration the uncertainty of measuring y, but assumed that we know x exactly. In real life, there is also some uncertainty in measuring x as well. How can we take this uncertainty into consideration?

For the problem of finding local minima and maxima, this uncertainty was taken into consideration in [9]. It is desirable to extend this approach to finding the range of the derivatives.

Parallelization. In the above text, we described how to compute the derivative of an interval-valued function in time $O(n^2)$, where n is the number of observations, and showed that this algorithm is (asymptotically) optimal in the sense that no algorithm can compute this derivative faster.

For reasonable n, e.g., for $n \approx 10^3$, n^2 computational steps means a million steps; it is quite doable on modern computers. However, for large n, e.g., for $n \approx 10^6$, n^2 computational steps is 10^{12} steps, so on a modern Gigaherz machine, the corresponding computations will take 10^3 sec – almost an hour.

How can we further speed up the corresponding computations? Our optimality result shows that we cannot achieve a drastic speed-up if we use sequential computers. Thus, the only way to speed up the corresponding computations is to use *parallel* computers.

For the problem of finding local minima and local maxima, parallel computers can indeed speed up the corresponding computations; see, e.g., [21]. An important question is therefore: How can speed up the computation of the corresponding derivative by using parallel computers?

Acknowledgments

This work was supported in part by NASA under cooperative agreement NCC5-209, by the Future Aerospace Science and Technology Program (FAST) Center for Structural Integrity of Aerospace Systems, effort sponsored by the Air Force Office of Scientific Research, Air Force Materiel Command, USAF, under grant F49620-00-1-0365, by NSF grants EAR-0112968 and EAR-0225670, by

the Army Research Laboratories grant DATM-05-02-C-0046, by the Hewlett-Packard equipment grants 89955.1 and 89955.2, by the Personal Interface AccessGrid awarded by the Education, Outreach and Training Partnership for Advanced Computational Infrastructure EOT-PACI, by the IEEE/ACM SC2003 Minority Serving Institutions Participation Grants, and by the Brazilian funding agencies CNPq/CTPETRO, CNPq/CTINFO, and FAPERGS.

The authors are thankful to participants of the interval computations mailing list, especially to R. Baker Kearfott, Svetoslav Markov, Arnold Neumaier, and Andrei Sobolevskii, for valuable comments.

References

[1] M. S. Aguiar, A. C. R. Costa, and G. P. Dimuro, "ICTM: an interval tessellation-based model for reliable topographic segmentation", *Numerical Algorithms* (to appear).

[2] R. Angelov and S. Markov, "Extended Segment Analysis", *Freiburger Intervall-Berichte*, 1981, Vol. 81(10), pp. 1–63.

[3] D. D. Coblentz, V. Kreinovich, B. S. Penn, and S. A. Starks, "Towards Reliable Sub-Division of Geological Areas: Interval Approach" *Proceedings of the 19th International Conference of the North American Fuzzy Information Society NAFIPS'2000*, Atlanta, Georgia, July 13–15, 2000, pp. 368–372.

[4] D. D. Coblentz, V. Kreinovich, B. S. Penn, and S. A. Starks, "Towards Reliable Sub-Division of Geological Areas: Interval Approach", In: L. Reznik and V. Kreinovich (eds.), *Soft Computing in Measurements and Information Acquisition*, Springer-Verlag, Berlin-Heidelberg, 2003, pp. 223–233.

[5] G. J. Deboeck, K. Villaverde, and V. Kreinovich, "Interval Methods for Presenting Performance of Financial Trading Systems", *Reliable Computing*, 1995, Supplement (Extended Abstracts of APIC'95: International Workshop on Applications of Interval Computations, El Paso, TX, Febr. 23–25, 1995), pp. 67–70.

[6] L. Jaulin, M. Keiffer, O. Didrit, and E. Walter, *Applied Interval Analysis*, Springer-Verlag, Berlin, 2001.

[7] R. B. Kearfott, *Rigorous Global Search: Continuous Problems*, Kluwer, Dordrecht, 1996.

[8] R. B. Kearfott and V. Kreinovich V. (eds.), *Applications of Interval Computations*, Kluwer, Dordrecht, 1996.

[9] J. Lorkowski and V. Kreinovich, "If we measure a number, we get an interval. What if we measure a function or an operator?", Reliable Computing, 1996, Vol. 2, No. 3, pp. 287–298.

[10] S. Markov, "Calculus for interval functions of a real variable", *Computing*, 1979, Vol. 22, pp. 325–337.

[11] R. E. Moore, *Methods and Applications of Interval Analysis*, SIAM, Philadelphia, 1979.

[12] H. Muñoz and R. B. Kearfott, "Slope intervals, generalized gradients, semigradients, and csets", *Reliable Computing* (to appear).

[13] A. Neumaier, *Introduction to Numerical Analysis*, Cambridge Univ. Press, Cambridge, 2001.

[14] H. T. Nguyen and V. Kreinovich, "How to Divide a Territory? A New Simple Differential Formalism for Optimization of Set Functions", *International Journal of Intelligent Systems*, 1999, Vol. 14, No. 3, pp. 223–251.

[15] D. Ratz, *Automatic Slope Computation and its Application in Nonsmooth Global Optimization*, Shaker-Verlag, Aachen, 1998.

[16] B. Sendov, "Segment arithmetic and segment limit", *C. R. Acad. Bulg. Sci.*, 1977, Vol. 30, pp. 955–968.

[17] B. Sendov, "Segment derivatives and Taylor's formula", *C. R. Acad. Bulg. Sci.*, 1977, Vol. 30, pp. 1093–1096.

[18] B. Sendov, "Some topics of segment analysis", In: K. Nickel (ed.), *Interval Mathematics'80*, Academic Press, N.Y., 1980, pp. 236–245.

[19] B. Sendov, *Hausdorff Approximations*, Kluwer, Dordrecht, 1990.

[20] K. Villaverde and V. Kreinovich, "A linear-time algorithm that locates local extrema of a function of one variable from interval measurement results," *Interval Computations*, 1993, No. 4, pp. 176–194.

[21] K. Villaverde and V. Kreinovich, "Parallel algorithm that locates local extrema of a function of one variable from interval measurement results", *Reliable Computing*, 1995, Supplement (Extended Abstracts of APIC'95: International Workshop on Applications of Interval Computations, El Paso, TX, Febr. 23–25, 1995), pp. 212–219.