

# Why Gaussian and Cauchy Functions Are Efficient in Filled Function Method: A Possible Explanation

José Guadalupe Flores Muñiz<sup>1</sup>, Vyacheslav V. Kalashnikov<sup>2,3</sup>,  
Nataliya Kalashnykova<sup>1,4</sup>, and Vladik Kreinovich<sup>5</sup>

<sup>1</sup>Department of Physics and Mathematics  
Universidad Autónoma de Nuevo León  
San Nicolás de los Garza, México  
jose\_guadalupe64@hotmail.com  
nkalash2009@gmail.com

<sup>2</sup>Department of Systems and Industrial Engineering  
Tecnológico de Monterrey ITESM, Campus Monterrey  
Monterrey, Mexico, kalash@itesm.mx

<sup>3</sup>Department of Experimental Economics  
Central Economics and Mathematics Institute (CEMI)  
Moscow, Russian Federation

<sup>4</sup>Department of Computer Science  
Sumy State University  
Sumy, Ukraine

<sup>5</sup>Department of Computer Science  
University of Texas at El Paso  
El Paso, Texas 79968, USA, vladik@utep.edu

One of the main problems of optimization algorithms is that they end up in a local optimum. It is therefore necessary to make sure that the algorithm gets out of the local optimum and eventually reaches the global optimum. One of the promising methods to leave the local optimum is the filled function method. It turns out that empirically, the best smoothing functions to use in this method are the Gaussian and the Cauchy functions. In this talk, we provide a possible theoretical explanation for this empirical result.