

Fair Economic Division: How to Modify Shapley Value to Take Into Account that Different People Have Different Productivity

Christian Servin¹ and Vladik Kreinovich²

¹El Paso Community College, cservin1@epcc.edu

²Department of Computer Science, University of Texas at El Paso, vladik@utep.edu

Formulation of the problem

Fair division: a problem. Let us assume that a group $N_n \stackrel{\text{def}}{=} \{1, \dots, n\}$ of n people jointly gets some benefit $v(N_n)$. What is the fair way to distribute this benefit between the participants, i.e., to assign values $\varphi_1, \dots, \varphi_n$ whose sum is $v(N_n)$?

What do we need to know to make a fair division. To make a fair distribution, it is important to know what is the contribution of each participant. This can be described by providing, for each subset $S \subseteq N$, a value $v(S)$ that people from S would have gained if they acted on their own, without help of others. So, we get a function $S \mapsto v(S)$ that characterizes the situation.

Shapley value. A Nobelist Lloyd Shapley found out that under reasonable conditions, there is only one way $\varphi_i(v)$ to assign the distribution to each such function $v(S)$; see, e.g., [?, ?, ?, ?, ?, ?]. This distribution has many equivalent forms, In this paper, we use the following form

$$\varphi_i(v) = \sum_{S: i \in S} \frac{t(S)}{|S|}, \quad (1)$$

where $|S|$ denotes the number of elements in the set S and

$$t(S) \stackrel{\text{def}}{=} \sum_{R \subseteq S} (-1)^{|S|-|R|} \cdot v(R). \quad (2)$$

What are the requirements behind Shapley value. Shapley's first condition is *symmetry*: if two participants i and j contribute equally, i.e., if the values $v(S)$ do not change when we swap i and j , then these participants should get equal amounts: $\varphi_i(v) = \varphi_j(v)$.

Shapley's second condition is that if a person i is not contributing, i.e., if $v(S \cup \{i\}) = v(S)$ for all S , then we should have $\varphi_i(v) = 0$.

Shapley's third condition is additivity: if we have two situations $u(S)$ and $v(S)$, then we can:

- either consider them separately
- or view them as a single situation with gain $w(S) = u(S) + v(S)$.

The outcome should not depend on how we view this, so we should have $\varphi_i(w) = \varphi_i(u) + \varphi_i(v)$.

Comment. In this paper, we will only deal with economic applications of Shapley value. It should be mentioned that Shapley value is now also actively used in machine learning, to find the importance $\varphi_i(v)$ of each of n features based on the effectiveness $v(S)$ of solving the problem when we only use features from the set S .

Why go beyond Shapley value. Symmetry makes perfect sense if all participants are equally productive. In reality, people have different productivity: e.g., some programmers are several times more productive than others. If we naively apply Shapley value to compute each person's bonus, more productive participants will get the exact same amount as less productive ones, which is not fair. It is therefore desirable to take productivity p_i of each participant into account. In other words, we need to determine the values φ_i based on both $v(S)$ and the values $p = (p_1, p_2, \dots)$: $\varphi_i(v, p)$.

What we do in this paper. In this paper, we show how to adjust the requirement behind the Shapley value so that they would lead to a unique determination of the desired distributions $\varphi_i(v, p)$.

Comment. One can easily see that if all participants have the same productivity, i.e., if $p_1 = \dots = p_n$, then the new formula (3) becomes the usual Shapley value formula (1).

What we propose

Natural first requirement. If i 's productivity is twice larger than j 's, this means that the company can replace i with two less productive workers and get the same result. After this replacement, all participants have the same productivity, so to this replaced situation, we can apply symmetry and get Shapley value – and then assign to i the sum of bonuses that Shapley value recommends for his/her two replacements.

Similarly, if we replace person i with 3 or more workers, it makes sense to require that the amount given to the person i should be equal to the sum of amounts given to these workers. It turns out that it is sufficient to require this property only for situation which are *jit simple* – in some precise sense described below.

Natural second requirement. If the productivity changes a little bit, the resulting distribution should also change a little bit. In mathematical terms, this means that the dependence of distribution on productivity should be continuous.

Main result. If impose these two additional requirements, then we get the following result.

Definition 1. By a situation, we mean a triple (n, v, p) , where:

- n is a positive integer,
- v is a function that assigns a value $v(S) \geq 0$ to each subset $S \subseteq N_n \stackrel{\text{def}}{=} \{1, \dots, n\}$ and for which $R \subseteq S$ implies $v(R) \leq v(S)$, and
- $p = (p_1, \dots, p_n)$ is a tuple of n positive numbers.

Definition 2. We say that the situation (n, v, p) is simple if for some subset $R \subseteq N_n$, we have $v(S) = 0$ if $R \not\subseteq S$ and $v(S) = v(R)$ otherwise. We will call this set R basic.

Definition 3.

- By a division strategy, we mean a function $\varphi(n, v, p)$ that assigns, to each situation (n, v, p) , an n -tuple of real numbers $\varphi_i(n, v, p)$, $1 \leq i \leq n$, for which $\varphi_1(n, v, p) + \dots + \varphi_n(n, v, p) = v(N_n)$.
- We say that a division strategy is symmetric if for every situation in which swapping i and j does not change v and p , i.e., in which $p_i = p_j$ and $v(S \cup \{i\}) = v(S \cup \{j\})$ for all sets S that contain neither i nor j , we have $\varphi_i(n, v, p) = \varphi_j(n, v, p)$.
- We say that a division strategy has the null-player property if for each situation and for each player i for which $v(S \cup \{i\}) = v(S)$ for all S , we get $\varphi_i(n, v, p) = 0$.
- We say that a division strategy is additive if for all u and v , we have $\varphi_i(n, u + v, p) = \varphi_i(n, u, p) + \varphi_i(n, v, p)$ for all i .
- We say that a division strategy is continuous if $\varphi(n, v, p)$ is a continuous function of p .
- We say that a division strategy is productivity-based if for every simple situation with a basic set R , if we combine participants from a subset $R' \subseteq R$ into a single participant i_0 with productivity equal to the sum of productivities of all members of R' , then in this new situation (n', v', p') ,

$$\varphi_{i_0}(n', v', p') = \sum_{i \in R'} v_i(n, v, p).$$

Proposition. There exists one and only one division strategy which is symmetric, has null-player property, is additive, continuous, and productivity-based. In this strategy,

$$\varphi(n, v, p) = p_i \cdot \sum_{S: i \in S} \frac{t(S)}{p(S)}, \quad (3)$$

where we denoted

$$p(S) \stackrel{\text{def}}{=} \sum_{i \in S} p_i. \quad (4)$$